Cochrane
Library

Cochrane Database of Systematic Reviews

Pharmacological treatment for Buerger's disease (Review)

Cacione DG, Macedo CR, Baptista-Silva JCC

Cacione DG, Macedo CR, Baptista-Silva JCC.

Pharmacological treatment for Buerger's disease.

Cochrane Database of Systematic Reviews 2016, Issue 3. Art. No.: CD011033.

DOI: 10.1002/14651858.CD011033.pub3.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	7
OBJECTIVES	8
METHODS	8
RESULTS	10
Figure 1.	11
Figure 2.	12
Figure 3.	13
Figure 4.	16
ADDITIONAL SUMMARY OF FINDINGS	17
DISCUSSION	24
AUTHORS' CONCLUSIONS	25
ACKNOWLEDGEMENTS	25
REFERENCES	26
CHARACTERISTICS OF STUDIES	28
DATA AND ANALYSES	37
Analysis 1.1. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 1 Ulcer healing (8 weeks) - iloprost 200 mcg.	38
Analysis 1.2. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 2 Ulcer healing (8 weeks) - iloprost 400 mcg.	39
Analysis 1.3. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 3 Ulcer healing (6 months) - iloprost 200 mcg.	39
Analysis 1.4. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 4 Ulcer healing (6 months) - iloprost 400 mcg.	40
Analysis 1.5. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 5 Complete relief of rest pain (8 weeks) - iloprost 200 mcg.	40
Analysis 1.6. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 6 Complete relief of rest pain (8 weeks) - iloprost 400 mcg.	41
Analysis 1.7. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 7 Complete relief of rest pain (6 months) - iloprost 200 mcg.	41
Analysis 1.8. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 8 Complete relief of rest pain (6 months) - iloprost 400 mcg.	42
Analysis 1.9. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 9 Rate of amputation - iloprost 200 mcg.	42
Analysis 1.10. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 10 Rate of amputation - iloprost 400 mcg.	43
Analysis 2.1. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 1 Ulcer healing (4 weeks).	43
Analysis 2.2. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 2 Ulcer healing (6 months).	44
Analysis 2.3. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 3 Complete relief of rest pain (4 weeks).	44
Analysis 2.4. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 4 Rate of amputation.	45
Analysis 3.1. Comparison 3 Prostacyclin versus prostaglandin E1, Outcome 1 Ulcer healing.	45
Analysis 3.2. Comparison 3 Prostacyclin versus prostaglandin E1, Outcome 2 Complete relief of pain.	46
Analysis 4.1. Comparison 4 Folic acid versus placebo, Outcome 1 Pain (0 month).	46
Analysis 4.2. Comparison 4 Folic acid versus placebo, Outcome 2 Pain (2 months).	47

Analysis 4.3. Comparison 4 Folic acid versus placebo, Outcome 3 Pain (6 months).	47
Analysis 4.4. Comparison 4 Folic acid versus placebo, Outcome 4 Change in rate of amputation (2 months).	48
Analysis 4.5. Comparison 4 Folic acid versus placebo, Outcome 5 Change in rate of amputation (6 months).	48
APPENDICES	48
FEEDBACK	51
WHAT'S NEW	52
CONTRIBUTIONS OF AUTHORS	53
DECLARATIONS OF INTEREST	53
SOURCES OF SUPPORT	53
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	54
INDEX TERMS	54

Pharmacological treatment for Buerger's disease

Daniel G Cacione¹, Cristiane R Macedo², Jose CC Baptista-Silva³

¹Department of Surgery, UNIFESP - Escola Paulista de Medicina, São Paulo, Brazil. ²Brazilian Cochrane Centre, Centro de Estudos de Saúde Baseada em Evidências e Avaliação Tecnológica em Saúde, São Paulo, Brazil. ³Evidence Based Medicine, Cochrane Brazil, Universidade Federal de São Paulo, São Paulo, Brazil

Contact address: Daniel G Cacione, Department of Surgery, UNIFESP - Escola Paulista de Medicina, Rua Borges Lagoa, 564 cj 124, Vila Clementino, São Paulo, 04038000, Brazil. dancac@gmail.com. cacione@gmail.com.

Editorial group: Cochrane Vascular Group.

Publication status and date: Edited (no change to conclusions), comment added to review, published in Issue 3, 2016.

Review content assessed as up-to-date: 13 April 2015.

Citation: Cacione DG, Macedo CR, Baptista-Silva JCC. Pharmacological treatment for Buerger's disease. *Cochrane Database of Systematic Reviews* 2016, Issue 3. Art. No.: CD011033. DOI: 10.1002/14651858.CD011033.pub3.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Buerger's disease (thromboangiitis obliterans) is a non-atherosclerotic, segmental inflammatory pathology that most commonly affects the small and medium sized arteries, veins, and nerves in the upper and lower extremities. The etiology is unknown, but involves hereditary susceptibility, tobacco exposure, immune and coagulation responses. In many cases, there is no possibility of revascularization to improve the condition. Pharmacological treatment is an option for patients with severe complications, such as ischaemic ulcers or rest pain.

Objectives

To assess the effectiveness of any pharmacological agent (intravenous or oral) compared with placebo or any other pharmacological agent in patients with Buerger's disease.

Search methods

The Cochrane Vascular Trials Search Co-ordinator searched their Specialised Register (last searched in April 2015) and the Cochrane Register of Studies (Issue 3, 2015). The review authors searched trial registers and the European grey literature; screened reference lists of relevant studies, and contacted study authors and major pharmaceutical companies.

Selection criteria

Randomised controlled trials (RCTs) involving pharmacological agents used in the treatment of Buerger's disease.

Data collection and analysis

Two review authors, independently assessed the studies, extracted data and performed data analysis.

Main results

Five randomised controlled trials (total 602 participants) compared prostacyclin analogue with placebo, aspirin, or a prostaglandin analogue, and folic acid with placebo. No studies assessed other pharmacological agents such as cilostazol, clopidogrel and pentoxifylline or compared oral versus intravenous prostanoid.

Compared with aspirin, intravenous prostacyclin analogue iloprost improved ulcer healing (risk ratio (RR) 2.65; 95% confidence interval (CI) 1.15 to 6.11; 98 participants; one study; moderate quality evidence), and helped to eradicate rest pain after 28 days

(RR 2.28; 95% CI 1.48 to 3.52; 133 participants; one study; moderate quality evidence), although amputation rates were similar six months after treatment (RR 0.32; 95% CI 0.09 to 1.15; 95 participants; one study; moderate quality evidence). When comparing prostacyclin (iloprost and clinprost) with prostaglandin (alprostadil) analogues, ulcer healing was similar (RR 1.13; 95% CI 0.76 to 1.69; 89 participants; two studies; $I^2 = 0\%$; very low quality evidence), as was the eradication of rest pain after 28 days (RR 1.57; 95% CI 0.72 to 3.44; 38 participants; one study; low quality evidence), while amputation rates were not measured. Compared with placebo, the effects of oral prostacyclin analogue iloprost were similar for: healing ischaemic ulcers (iloprost 200 mcg; RR 1.11; 95% CI 0.54 to 2.29; 133 participants; one study; moderate quality evidence, and iloprost 400 mcg; RR 0.90; 95% CI 0.42 to 1.93; 135 participants; one study; moderate quality evidence), eradication of rest pain after eight weeks (iloprost 200 mcg; RR 1.14; 95% CI 0.79 to 1.63; 207 participants; one study; moderate quality evidence, and iloprost 400 mcg; RR 1.11; 95% CI 0.77 to 1.59; 201 participants; one study; moderate quality evidence), and amputation rates after six months (iloprost 200 mcg; RR 0.54; 95% CI 0.19 to 1.56; 209 participants; one study, and iloprost 400 mcg; RR 0.42; 95% CI 0.13 to 1.31; 213 participants; one study). When comparing folic acid with placebo in patients with Buerger's disease and hyperhomocysteinaemia, pain scores were similar, there were no new cases of amputation in either group, and ulcer healing was not assessed (very low quality evidence).

Treatment side effects such as headaches, flushing or nausea were not associated with treatment interruptions or more serious consequences. Outcomes such as amputation-free survival, walking distance or pain-free walking distance, and ankle brachial index were not assessed by any study.

Overall, the quality of the evidence was very low to moderate, with few studies, small numbers of participants, variation in severity of disease of participants between studies and missing information regarding for example baseline tobacco exposure.

Authors' conclusions

Moderate quality evidence suggests that intravenous iloprost (prostacyclin analogue) is more effective than aspirin for eradicating rest pain and healing ischaemic ulcers in Buerger's disease, but oral iloprost is not more effective than placebo. Verylow and low quality evidence suggests there is no difference between prostacyclin (iloprost and clinprost) and the prostaglandin analogue alprostadil for healing ulcers and relieving pain respectively in severe Buerger's disease. Very-low quality evidence suggests there is no difference in pain scores and amputation rates between folic acid and placebo, in people with Buerger's disease and hyperhomocysteinaemia. High quality trials assessing the effectiveness of pharmacological agents (intravenous or oral) in people with Buerger's disease are needed.

PLAIN LANGUAGE SUMMARY

Pharmacological treatment for Buerger's disease

Background

Buerger's disease is characterized by recurring progressive inflammation and clotting in small and medium arteries and veins of the hands and feet. Its cause is unknown, but it is most common in men with a history of tobacco use. It is responsible for ulcers and extreme pain in the limbs of young smokers. In many cases, mainly in patients with the most severe form, there is no possibility of improving the condition with surgery, and therefore, drugs (pharmacological agents) are used. These can be pharmacological agents, such as cilostazol, clopidogrel, and pentoxifylline, or medicine derivatives of prostacyclin and prostaglandin, which redirect blood flow and improve the circulation in affected areas, and theoretically, help to heal ulcers and relieve rest pain. This review assessed the effectiveness of pharmacological agents in the treatment of patients with Buerger's disease.

Key results

Our search identified five randomised controlled trials, with a total of 602 participants and a treatment period of around four weeks (evidence current until April 2015). The comparisons included prostacyclin analogue versus placebo, aspirin, and a prostaglandin analogue, and folic acid versus placebo. We did not identify studies that assessed pharmacological agents such as cilostazol, clopidogrel and pentoxifylline, or studies that compared oral prostanoid versus intravenous prostanoid. The included studies assessed derivatives of prostacyclin and prostaglandin, which have the ability to redirect blood flow and improve the circulation in affected areas.

Moderate quality evidence from one study suggested that intravenous iloprost was effective in healing ulcers and relieving rest pain after 28 days of treatment when compared with oral aspirin, but no differences were found in the rates of amputation. Evidence from two studies suggested that prostacyclin was as effective as prostaglandin analogues in healing ulcers (very low quality evidence) and eradicating pain at rest (low quality evidence), but rates of amputation were not assessed. Moderate quality evidence from one study suggested

that there was no difference between placebo and the oral prostacyclin analogue iloprost (200 mcg and 400 mcg) in healing ischaemic ulcers or eradicating pain at rest after eight weeks and six months, and rates of amputation after six months. Very-low quality evidence from one study showed no difference between placebo and folic acid, in patients with Buerger's disease and hyperhomocysteinaemia (a medical condition characterized by abnormally high level of homocysteine in the blood), in rates of amputation and pain scores. Ulcer healing was not measured. Treatment side effects, such as headaches or nausea, did not result in treatment interruptions or more serious consequences. Outcomes such as amputation-free survival, walking distance or pain-free walking distance, and ankle brachial index were not assessed by any study.

Quality of the evidence

Overall, the quality of the evidence was very low to moderate, with few studies, small numbers of participants, variation in severity of disease of participants between studies and missing information regarding for example baseline tobacco exposure. High quality trials assessing the effectiveness of pharmacological agents (intravenous or oral) in people with Buerger's disease are needed.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Oral prostacyclin analogue (iloprost) versus placebo for treatment of Buerger's disease						
Patient or population: patients with Buerger's disease Settings: hospital and community Intervention: oral prostacyclin analogue (iloprost) in two doses: 200 mcg and 400 mcg Comparison: placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	placebo	oral prostacyclin analogue				
Ulcer healing (200 mcg) Follow-up: 8 weeks	Study population		RR 1.11 (0.54 to 2.29)	133 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	171 per 1000	190 per 1000 (93 to 393)				
Ulcer healing (400 mcg) Follow-up: 8 weeks	Study population		RR 0.90 (0.42 to 1.93)	135 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	171 per 1000	154 per 1000 (72 to 331)				
Ulcer healing (200 mcg) Follow-up: 6 months	Study population		RR 1.19 (0.82 to 1.73)	133 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	414 per 1000	493 per 1000 (340 to 717)				
Ulcer healing (400 mcg) Follow-up: 6 months	Study population		RR 1.00 (0.67 to 1.50)	135 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	414 per 1000	414 per 1000 (278 to 621)				

Complete relief of rest pain (200 mcg) Follow-up: 8 weeks	Study population		RR 1.14 (0.79 to 1.63)	207 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate
	343 per 1000	391 per 1000 (271 to 559)			
Complete relief of rest pain (400 mcg) Follow-up: 8 weeks	Study population		RR 1.11 (0.77 to 1.59)	210 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate
	343 per 1000	381 per 1000 (264 to 546)			
Complete relief of rest pain (200 mcg) Follow-up: 6 months	Study population		RR 1.28 (1.00 to 1.64)	207 (1 RCT)	⊕⊕○○ ^{1,2,3} low
	490 per 1000	627 per 1000 (490 to 804)			
Complete relief of rest pain (400 mcg) Follow-up: 6 months	Study population		RR 1.00 (0.76 to 1.32)	210 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate
	490 per 1000	490 per 1000 (373 to 647)			
Rate of amputation (200 mcg) Follow-up: 6 months	Study population		RR 0.54 (0.19 to 1.56)	209 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate
	87 per 1000	47 per 1000 (17 to 136)			
Rate of amputation (400 mcg) Follow-up: 6 months	Study population		RR 0.42 (0.13 to 1.31)	213 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate
	87 per 1000	37 per 1000 (11 to 114)			

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk Ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

- ¹ one single study (doubt about reproducibility of data), downgraded by one level
- ² conflict of interest not stated but it was not considered sufficient to downgrade the quality of evidence
- ³ inconsistent with dose-response effect, downgraded by one level

BACKGROUND

Description of the condition

Buerger's disease (thromboangiitis obliterans) is a non-atherosclerotic, segmental inflammatory pathology that most commonly affects the small and medium sized arteries, veins and nerves in the upper and lower extremities (Olin 2000). Von Winiwarter first described a patient with the disease in 1879 (von Winiwarter 1879), but it was Leo Buerger, in 1908, who published a detailed description of the pathological findings on amputated limbs and named the disease (Buerger 1908).

The prevalence of the disease among all patients with peripheral arterial disease (PAD) varies from as low as 0.5% to 5.6% in Western Europe to as high as 45% to 63% in India, and 16% to 66% in Korea and Japan (Cachovan 1988; Malecki 2009; Olin 2000). The etiology is unknown, but involves hereditary susceptibility, tobacco exposure, immune and coagulation responses (Malecki 2009). Currently, a possible infectious etiology is gaining interest, especially after the findings of micro-organisms of the oral flora in occlusive thrombi in patients with Buerger's disease and moderate to severe periodontitis (Iwai 2005; Li 2008). Another hypothesis is the possible pathogenic role of rickettsial infection in Buerger's disease (Bartolo 1987; Fazeli 2013). Features distinguishing Buerger's disease from atherosclerosis include the pathology distribution (with involvement of both the upper and lower extremities), associated superficial venous thrombosis, a paucity of atherosclerotic risk factors and normal proximal large arteries (Weinberg 2012).

Diagnosis and complications

The typical patient with Buerger's disease is a young man (younger than 40 or 45 years) with a history of tobacco use, who presents with progressive claudication, ischaemic ulcers, or pain at rest (Olin 2000); approximately 76% of patients have ischaemic ulcerations at the time of presentation (Olin 2006). To date, there are no unanimous diagnostic criteria for Buerger's disease. The most commonly used are Shionoya's criteria, which comprise: (1) smoking history; (2) onset of symptoms before the age of 50 years; (3) infrapopliteal arterial occlusions; (4) either arm involvement or phlebitis migrans; and (5) absence of atherosclerotic risk factors other than smoking (Shionoya 1983). All criteria should be present. The disease is usually confined to the distal circulation and is almost always infrapopliteal in the legs and distal to the brachial artery in the arms (Olin 2000). In fact, the distal and diffuse nature of the disease culminates in critical limb ischaemia (CLI) in approximately 76% to 81% of patients, with poor chances of revascularization (Olin 2006). In patients diagnosed with limb ischaemia, as in Buerger's disease, the clinical evaluation is done according to the Rutherford classification and the Fontaine stages. The Rutherford classification for PAD has seven categories. These are: (0)

asymptomatic; (1) mild claudication; (2) moderate claudication; (3) severe claudication; (4) rest pain; (5) minor tissue loss, non-healing ulcer or focal gangrene with diffuse pedal ischaemia; (6) major tissue loss extending above the transmetatarsal level; and (7) a functional foot that is no longer salvageable (Rutherford 2005). The Fontaine classification has four stages: (I) asymptomatic; (II) intermittent claudication (IC); (III) rest pain; (IV) ulceration or gangrene, or both (Fontaine 1954). Novo 2004 described a modified Fontaine classification, the Leriche-Fontaine classification: (I) asymptomatic or effort pain; (IIA) effort pain or pain-free walking distance further than 200 metres; (IIB) pain-free walking distance less than 200 metres; (IIIA) rest pain, ankle arterial pressure higher than 50 mm Hg; (IIIB) rest pain, ankle arterial pressure lower than 50 mm Hg; (IV) trophic lesions, necrosis or gangrene (Novo 2004). According to Cooper 2004, the risk of any extremity amputation during 15.6 years of follow-up in patients with Buerger's disease is 25% at five years, 38% at 10 years and 46% at 20 years.

Description of the intervention

In patients with CLI and poor chances of revascularization, as seen in many patients diagnosed with Buerger's disease, pharmacological treatment is given to improve the blood flow (perfusion) in the affected extremity. The most commonly used pharmacological agents are aspirin, cilostazol (Bedenis 2014; Ruffolo 2010), prostanooids (Malecki 2009), and bosentan (De Haro 2009).

Aspirin is a drug that inhibits cyclo-oxygenase, which is responsible for the synthesis of thromboxane and prostaglandins. It is important in cardiac and cerebrovascular atherosclerotic diseases, as it inhibits platelet aggregation. The most important contraindications are hypersensitivity to salicylates, active gastrointestinal ulcers, patients with hemorrhagic disorders, renal and hepatic failure, pregnancy and use in children. Aspirin is administered orally with a recommended dosage of 75 to 325 mg (Brunton 2011).

Cilostazol is a quinolinone derivative drug that inhibits cellular phosphodiesterase (more specific for phosphodiesterase III), affecting both vascular beds and cardiovascular function. It causes a non-homogeneous dilation of vascular beds, with greater dilation in the femoral beds than in vertebral, carotid or superior mesenteric arteries. In other words, cilostazol 'steals' a small part of the blood from other territories (gastrointestinal and cerebral) to improve perfusion in ischaemic limbs. A further action of cilostazol is the reversible inhibition of platelet aggregation. Cilostazol is contraindicated in patients with congestive heart failure (CHF) of any severity, haemostatic disorders or active pathologic bleeding, such as bleeding peptic ulcer and intracranial bleeding, and in patients with known or suspected hypersensitivity to cilostazol. The more common side effects of cilostazol use include headaches, diarrhoea, abnormal stools and palpitations (Chapman 2003). Cilostazol is administered orally and is available in 50 mg or 100 mg tablets. Cilostazol was approved by the US Food and Drug Administration (FDA) in 1999 for the reduction of symptoms of intermit-

tent claudication as a result of atherosclerosis (Dindyal 2009; FDA 1999).

Prostanoids (prostaglandin and prostacyclin analogues) are eicosanoid derivatives, commonly used for many conditions including pulmonary hypertension, sexual impotence, and glaucoma. Because of their short half-life, around two to three minutes, these synthetic drugs must be administered by continuous intravenous infusion (Safdar 2011). The development of stable prostacyclin analogues (such as iloprost) with a longer half-life has allowed the oral use of these drugs. The most important contraindications are heart failure (caused by arrhythmias, myocardial pathology, valvulopathy, or coronary insufficiency), intracranial haemorrhage, gastrointestinal ulcers, and trauma. Side effects include headache, flushing, malaise, gastrointestinal distress and, with higher doses, hypotension. The maximum iloprost dose that is administered is around 2 ng/kg/min by continuous infusion (Grant 1992).

Bosentan is a potent and mixed endothelin-A and endothelin-B receptor antagonist, causing selective vasodilatory effects (Weber 1996). Some important reported side effects are hepatotoxicity and fluid retention. Bosentan is administered orally, mainly in patients with pulmonary arterial hypertension, with a recommended dosage of 62.5 mg (twice daily) to 125 mg (twice daily).

How the intervention might work

Aspirin (acetylsalicylic acid), a thromboxane production inhibitor, is well-known as an antiplatelet drug, and is used in heart attack and stroke prevention (Brunton 2011).

Cilostazol, a selective inhibitor of phosphodiesterase III, is used mainly in patients with IC, and acts as a direct arterial vasodilator and inhibits platelet aggregation (Rutherford 2005).

Prostanoids act by binding to specific receptors in the endothelium (causing vasodilatation) and platelets (inhibiting platelet aggregation), which causes a transitory increase in arterial peripheral perfusion (Brunton 2011). Arterial vasodilatation in ischaemic areas increases blood perfusion and, consequently, increases the chances of ulcer healing and improving rest pain. Inhibiting platelet aggregation prevents the occlusion of small arteries and, therefore, stabilizes the disease.

Bosentan, a potent and mixed endothelin-A and endothelin-B receptor antagonist, causes selective vasodilatory effects (Weber 1996). Bosentan has been used with success in patients with digital ulcers and systemic sclerosis (Launay 2006; Matucci-Cerinic 2011), opening the perspective for use in other conditions such as Buerger's disease (De Haro 2009).

Therefore, these pharmacological agents are used to improve perfusion in ischaemic limbs due to their vasodilatory and antiplatelet effects. The effects may vary from agent to agent.

Why it is important to do this review

Buerger's disease is a debilitating condition which can affect active, young people. In many cases, no possibility exists of revascularization to improve the condition. Different combinations of drugs, doses and administration pathways (oral and intravenously) have been approved for use. However, to date there is no consensus about the best pharmacological treatment for patients with Buerger's disease. A systematic review is opportune and extremely relevant.

OBJECTIVES

To assess the effectiveness of any pharmacological agent (intravenous or oral) compared with placebo or any other pharmacological agent in patients with Buerger's disease.

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials (RCTs) involving pharmacological agents used in the treatment of Buerger's disease.

Types of participants

Patients clinically diagnosed with Buerger's disease.

Types of interventions

We assessed any pharmacological agents used in treating patients with Buerger's disease, including drugs utilized for atherosclerotic diseases, such as aspirin or cilostazol, resulting in the possible comparisons listed below:

- (1) (Oral or intravenous) prostanoid (e.g. iloprost) versus placebo
- (2) Oral prostanoid versus intravenous prostanoid
- (3) (Oral or intravenous) prostanoid (e.g. iloprost) versus aspirin
- (4) (Oral or intravenous) prostanoid (e.g. iloprost) versus cilostazol
- (5) Aspirin versus placebo
- (6) Cilostazol versus placebo
- (7) Aspirin versus cilostazol
- (8) Any pharmacological agent versus placebo or any other pharmacological agent

Prostanoids could be either prostaglandin or prostacyclin analogues.

We excluded studies that did not assess pharmacological agents in the treatment of Buerger's disease.

Types of outcome measures

Primary outcomes

- (1) Ulcer healing
- (2) Pain: assessed using a validated pain score or scale, or quality of life (QoL) questionnaire
- (3) Rate of amputation: major amputation (defined as amputation of the lower or upper limb above the ankle or the wrist, respectively); and minor amputation (defined as amputation of a hand or foot or any part of)

Secondary outcomes

- (1) Amputation-free survival
- (2) Side effects of pharmacological agents, including bleeding, headache, flushing, or nausea
- (3) Walking distance or pain-free walking
- (4) Ankle brachial index

Search methods for identification of studies

There were no language restrictions.

Electronic searches

The Cochrane Vascular Trials Search Co-ordinator (TSC) searched their Specialised Register (last searched in April 2015) and the Cochrane Register of Studies (CRS) <http://www.metaxis.com/CRSWeb/Index.asp> (Issue 3, 2015). See [Appendix 1](#) for details of the search strategy used to search the CRS. The Specialised Register is maintained by the TSC and is constructed from weekly electronic searches of MEDLINE, EMBASE, CINAHL, AMED, and through handsearching relevant journals. The full list of the databases, journals and conference proceedings which have been searched, as well as the search strategies used, are described in the [Specialised Register](#) section of the Cochrane Vascular module in *The Cochrane Library* (www.cochranelibrary.com).

Trial registers

We searched the following trial registers:

- ISRCTN register (www.isrctn.com) on May 10, 2015;
- ClinicalTrials.gov (www.clinicaltrials.gov) on May 10, 2015;
- The Australian New Zealand Clinical Trials Registry (www.anzctr.org.au) on May 10, 2015;
- The EU Clinical Trials Register (www.clinicaltrialsregister.eu) on May 10, 2015;
- The World Health Organization International Clinical Trials Registry Platform (www.who.int/trialsearch) on May 10, 2015;

- LILACS, Latin American and Caribbean Health Sciences Literature (<http://lilacs.bvsalud.org/>) on May 10, 2015.

We used the terms 'Buerger's disease'; 'thromboangiitis obliterans'; 'von Winiwarter disease'; and word variations to perform the search.

Grey literature

We searched the grey literature produced in Europe by consulting the OpenGrey Database (www.opengrey.eu). We used the terms 'Buerger's disease'; 'thromboangiitis obliterans'; 'von Winiwarter disease'; and word variations to perform the search.

Searching other resources

We searched the reference lists of relevant articles retrieved by the electronic searches for additional citations.

We contacted study authors to inquire about ongoing or unpublished studies (Dr JN Fiessinger, Dr K Esato, Dr K Ishitobi, Dr M Verstraete, Dr MA Hoghoughi, and Dr S Ishimaru). Only Dr MA Hoghoughi replied, denying new studies. We asked pharmaceutical companies to provide information on both published and unpublished trials (Actelion Pharmaceuticals, AstraZeneca, Bayer Schering Pharma AG, Changzhou Highassay Chemical Co Ltd, LGM Pharma, Manus Aktteva Biopharma LLP, Otsuka Pharmaceutical Co Ltd, Pfizer Inc, Sandoz Inc). However, only AstraZeneca and Manus Aktteva Biopharma LLP replied to our correspondence, and both denied ongoing or unpublished studies.

Data collection and analysis

Selection of studies

Two review authors (DGC and JCCBS) independently assessed all studies that were identified by the search strategy for inclusion. Disagreements were resolved by the third author (CM).

Data extraction and management

For all eligible studies, two review authors (DGC and JCCBS) extracted data using the Cochrane Vascular's data extraction table. Where there were discrepancies, the third author (CM) solved disagreement. We entered the data into Review Manager 5.3.

Assessment of risk of bias in included studies

Two review authors (DGC and JCCBS) independently assessed the included studies for risk of bias using Cochrane's 'Risk of bias' tool as described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). The information about the risk of bias of the included studies was presented in the form of a table and a graph.

Measures of treatment effect

Dichotomous (categorical) data

Results were presented as summary risk ratios (RR) with 95% confidence intervals (CIs).

Continuous data

We had planned to use the mean difference (MD) with 95% CI where there was consistency in the outcome measure, or the standardised mean difference (SMD) to combine trials that measured the same outcome but used different methods.

Time-to-event data

We had planned to use hazard ratios (HR) with 95% CIs to measure the treatment effect for any time-to-event outcomes.

Unit of analysis issues

We considered the individual participant as the unit of randomisation for a single intervention.

Dealing with missing data

We contacted contact authors of included studies about methodological queries, but none of the contact authors answered the solicitation. Where possible, we had planned to analyse all outcome measures on an intention-to-treat basis by including data from all participants.

Assessment of heterogeneity

The heterogeneity among the eligible studies was quantified using the Chi² test and I² statistic, specifically using the formula $I^2 = (Q - df/Q) \times 100\%$ where Q was the Chi² statistic and df represented the degree of freedom. The I² statistic values were interpreted as follows:

- 0% to 25% = low heterogeneity;
- 25% to 75% = moderate heterogeneity;
- more than 75% = substantial heterogeneity (Higgins 2011)

Where substantial heterogeneity was detected, according to the criteria above, we had planned to perform a further investigation based on the pre-specified subgroup analysis.

Assessment of reporting biases

Had sufficient eligible trials been available, we had planned to explore publication bias through the use of funnel plots and to explore the presence of time-lag bias in both published and unpublished trials. As a result of the small number of included studies (five RCTs), these analyses were not performed.

Data synthesis

Review Manager (RevMan 5.3) was used to perform data synthesis. After assessing heterogeneity, we had planned to use a random-effects model meta-analysis if substantial heterogeneity between studies was detected. A fixed-effect model meta-analysis was planned if the studies estimated the same intervention effect and had low or moderate heterogeneity (Higgins 2011).

Subgroup analysis and investigation of heterogeneity

If sufficient information had been available, we had intended to perform subgroup analyses according to the following features:

- tobacco exposure (cigarette, cannabis, or any other form of smoking, either measured in a laboratory or declared) after the intervention;
- severity of the ischaemia, according to the Fontaine or Rutherford classification;
- different doses of the pharmacological agents.

Sensitivity analysis

We had intended to perform sensitivity analyses by looking separately at sponsored studies and publication bias, and by excluding studies with low and moderate methodological quality according to the 'Risk of bias' judgements. As a result of the small number of studies included for each comparison and a meta-analysis with only two studies, sensitivity analyses were not performed.

Summary of Findings

We presented the main findings of the review results for the quality of evidence, the magnitude of effect of the interventions examined, and the sum of available data on the primary outcomes (see [Types of outcome measures](#)) in 'Summary of findings' tables, according to Higgins 2011 and the GRADE Working group (Atkins 2004). Since we assessed different intervention comparisons, a 'Summary of findings' table was developed for each comparison included in the Results sections. The GRADEprofiler software was used to assist in the preparation of the 'Summary of findings' tables.

RESULTS

Description of studies

See [Characteristics of included studies](#), [Characteristics of excluded studies](#) and [Characteristics of studies awaiting classification](#) tables.

Results of the search

A flow diagram of the search results is shown in [Figure 1](#).

Figure 1. Study flow diagram.

Included studies

Five randomised controlled studies were included in this review, with a combined total of 602 participants (Beigi 2014; Esato 1995; Fiessinger 1990; Ishitobi 1991; Verstraete 1998).

Beigi 2014 reported on a study of 30 participants with Buerger's disease (Fontaine II of ischaemia) and hyperhomocysteinaemia (a well known and established risk factor for limb ischaemia in patients with atherosclerosis), who were receiving folic acid or placebo (single dose); they compared the number of major and minor amputations and pain in both groups.

In Esato 1995, 46 participants with Buerger's disease received a prostacyclin analogue (clinprost) or a prostaglandin analogue (alprostadil) for four weeks; they compared improvements in ischaemic ulcers and rest pain.

Fiessinger 1990 reported on a European multicentre randomised study that included 152 participants with Buerger's disease, in critical limb ischaemia (rest pain, ulcers or gangrene), who were receiving prostacyclin analogue (iloprost endovenously) or aspirin orally for 28 days.

Ishitobi 1991 was a trial of 134 participants with critical limb ischaemia, which included 55 participants diagnosed with Buerger's disease. The pharmacological treatment assessed the efficacy between a prostaglandin analogue (alprostadil) and a prostacyclin analogue (iloprost), both of them administered intravenously for 28 days.

Verstraete 1998 described a large multicentre randomised trial

with 319 participants with Buerger's disease with rest pain, ischaemic ulcers, or both, who were administered a prostaglandin analogue (iloprost) orally or placebo for eight weeks, with a six-month follow-up.

No studies were identified that compared oral prostanoid and intravenous prostanoid, (oral or intravenous) prostanoid (e.g. iloprost) and cilostazol, aspirin and placebo, cilostazol and placebo, and aspirin and cilostazol.

Excluded studies

Nine studies were excluded (Bozkurt 2006; Coscia 1972; He 2007; Musial 1986; Reichert 1975; Steinorth 1967; Sun 1993; Yang 2005; Zelikovsky 1973); the reasons for exclusion are described in the [Characteristics of excluded studies](#) table. In brief, the main reasons for exclusion were interventions with no pharmacological agents (e.g. acupuncture, sympathectomy; Bozkurt 2006; He 2007; Yang 2005), studies without patients with Buerger's disease (Steinorth 1967), or participants with mixed diagnoses without a separate description of the outcomes for participants with Buerger's disease (Coscia 1972; Zelikovsky 1973).

Risk of bias in included studies

See [Figure 2](#) and [Figure 3](#) for a graphical summary of methodological quality for the included studies, based on the 'Risk of bias' domains.

Figure 2. 'Risk of bias' graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. 'Risk of bias' summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Beigi 2014	+	+	+	+	+	-	-
Esato 1995	?	?	+	+	-	+	-
Fiessinger 1990	?	+	+	+	+	+	+
Ishitobi 1991	?	?	+	?	-	+	-
Verstraete 1998	?	?	+	?	+	+	?

Allocation

Four included studies (Esato 1995; Fiessinger 1990; Ishitobi 1991; Verstraete 1998) did not describe the method of randomisation and therefore, were classed as unclear risk of bias. Only Beigi 2014, after contact with study authors, was classed as low risk of bias because the trialists used a computerised randomisation method. Allocation concealment was only reported in the Fiessinger 1990 study, which reported the utilization of a centre of randomisation. Beigi 2014, after contact with study authors, described a method of allocation concealment based on computerised codes that were in the possession of a third person (not involved in drug administration or outcome assessment). The remaining three studies (Esato 1995; Ishitobi 1991; Verstraete 1998) were classified as unclear risk of bias.

Blinding

All five included studies were blinded for performance bias. Beigi 2014, Esato 1995 and Fiessinger 1990 were blinded for detection bias; however, Ishitobi 1991 and Verstraete 1998 did not report whether the outcome evaluators were blinded.

Incomplete outcome data

Beigi 2014 described a study without losses to follow-up. Follow-up was available to six months.

Esato 1995 reported they used intention-to-treat analyses, but in practice, adopted 'as-treated' (per protocol) analyses. All exclusions were explained. Follow-up was available to four weeks.

Fiessinger 1990 reported by intention-to-treat, and justified all the post randomisation exclusions. Follow-up was available to six months. Fiessinger 1990 was therefore judged to be at low risk of attrition bias.

In Ishitobi 1991, the authors did not clearly describe the losses to follow-up, declaring only that there were more losses in the iloprost group, but not reporting how many participants and possible reasons. Follow-up was available to four weeks. Ishitobi 1991 was judged to be of high risk of attrition bias.

Verstraete 1998 also reported by intention-to-treat, and justified all the post randomisation exclusions. Follow-up was available to six months. Therefore, Verstraete 1998 was judged to be at low risk of attrition bias.

Selective reporting

Esato 1995, Fiessinger 1990, Ishitobi 1991, and Verstraete 1998 described all outcomes and were judged to be at low risk of reporting bias. Beigi 2014 did not describe the presence of rest pain or ischaemic ulcers and was judged to be at high risk of reporting bias.

Other potential sources of bias

In Esato 1995 and Ishitobi 1991, the study authors did not describe tobacco exposure before and after the treatment and both were judged to be at high risk of other bias. Beigi 2014 was judged to be at high risk of other bias because participants without critical ischaemia were eligible, resulting in low chances to be amputated. Verstraete 1998 was judged to be at unclear risk of bias because there was no information about any potential conflict of interest. Fiessinger 1990 was judged to be at low risk of bias because the proportion of smokers and non-smokers in the study arms at the beginning and after treatment were described.

Effects of interventions

See: [Summary of findings for the main comparison](#) Summary of findings: Oral prostacyclin analogue (iloprost) versus placebo; [Summary of findings 2](#) Summary of findings: Intravenous prostacyclin analogue (iloprost) versus oral aspirin; [Summary of findings 3](#) Summary of findings: Intravenous prostacyclin analogue versus intravenous prostaglandin analogue; [Summary of findings 4](#) Summary of findings: Folic acid versus placebo

(Oral or intravenous) prostanoid versus placebo

Oral prostacyclin analogue (iloprost) versus placebo

One study assessed this comparison (Verstraete 1998).

Primary outcomes

(1) Ulcer healing: assessed at the end of treatment (eight weeks) and six months after the start of treatment. After eight weeks (end of treatment), complete healing of all ulcers was 19% in the 200 mcg iloprost group (12/63 participants; RR 1.11; 95% CI 0.54 to 2.29; $P = 0.78$), 15% in the 400 mcg iloprost group (10/65 participants; RR 0.90; 95% CI 0.42 to 1.93; $P = 0.78$), and 17% in the placebo group (12/70 participants; [Analysis 1.1](#); [Analysis 1.2](#)). After six months, the 200 mcg iloprost group reported complete healing of all ulcers in 49% (31/63 participants; RR 1.19; 95% CI 0.82 to 1.73; $P = 0.37$), the 400 mcg iloprost group reported 41% (27/65 participants; RR 1.00; 95% CI 0.67 to 1.50; $P = 0.99$), and the placebo group reported 41% (29/70 participants) complete healing of all ulcers ([Analysis 1.3](#); [Analysis 1.4](#)). None of these findings were statistically significant.

Verstraete 1998 also reported improvement of the most important ulcer (improved healing was not a defined outcome in this review): improvement of the most important ulcer was 55% in the 200 mcg iloprost group ($P = 0.056$ versus placebo), 63% in

the 400 mcg iloprost group ($P = 0.008$ versus placebo), and 40% in the placebo group after eight weeks (end of treatment). After six months, the 200 mcg iloprost group reported improvement of the most important ulcer in 84% ($P = 0.007$ versus placebo), the 400 mcg iloprost group reported improvement in 68% ($P = 0.297$ versus placebo), and the placebo group reported improvement in 62%. These findings were not statistically significant, except for 400 mcg iloprost at eight weeks of treatment and 200 mcg iloprost at six months when compared with placebo.

(2) Pain: assessed at the end of treatment (eight weeks) and six months after the start of treatment. After eight weeks (end of treatment), complete relief of rest pain was 39% in the 200 mcg iloprost group (41/105 participants; RR 1.14; 95% CI 0.79 to 1.63; $P = 0.48$), 38% in the 400 mcg iloprost group (41/108 participants; RR 1.11; 95% CI 0.77 to 1.59; $P = 0.58$), and 34% in the placebo group (35/102 participants; [Analysis 1.5](#); [Analysis 1.6](#)). After six months, the 200 mcg iloprost group reported complete relief of rest pain in 63% (66/105 participants; RR 1.28; 95% CI 1.00 to 1.64; $P = 0.05$), the 400 mcg iloprost group reported 49% (53/108 participants; RR 1.00; 95% CI 0.76 to 1.32; $P = 0.99$), and the placebo group reported 49% (50/102 participants) complete relief of rest pain ([Analysis 1.7](#); [Analysis 1.8](#)). None of these findings were statistically significant, except for the iloprost 200 mcg group after six months of treatment.

(3) Rate of amputation: Assessed at six months after the start of treatment, the placebo group reported 9% major amputations (9/103 participants), the 200 mcg iloprost group reported 4.7% major amputations (5/106 participants; RR 0.54; 95% CI 0.19 to 1.56; $P = 0.25$; [Analysis 1.9](#)), and the 400 mcg iloprost group reported 3.6% major amputations (4/110 participants; RR 0.42; 95% CI 0.13 to 1.31; $P = 0.13$; [Analysis 1.10](#)). These findings were not statistically significant.

Secondary outcomes

Side effects: Participants in the 400 mcg iloprost group developed about 25% more (68%) side effects than participants in the 200 mcg iloprost group (43%) and the placebo group (38%). According to the study authors, the most frequent side effect reported was headache (more than 20%), followed by vasodilatation and trismus (the latter two cases for the high-dose iloprost group only). Amputation-free survival, walking distance or pain-free walking, and ankle brachial index were not assessed by [Verstraete 1998](#).

(Oral or intravenous) prostanoid (e.g. iloprost) versus aspirin

Intravenous prostacyclin analogue (iloprost) versus oral aspirin

One study assessed this comparison ([Fiessinger 1990](#)).

Primary outcomes

(1) Ulcer healing: assessed at the end of treatment (day 28) and six months after the start of treatment. Complete healing of all ulcers, assessed by an independent evaluator, in the iloprost group was 35% (18/52 patients) compared to 13% (6/46 patients) in the aspirin group at day 28 (RR 2.65; 95% CI 1.15 to 6.11; $P = 0.02$; [Analysis 2.1](#)); after six months, complete healing of all ulcers in the iloprost group was 27.45% (14/51 participants) compared to 6.8% (3/44 participants) in the aspirin group (RR 4.03; 95% CI 1.24 to 13.10; $P = 0.02$; [Analysis 2.2](#)).

(2) Pain: assessed at the end of treatment (day 28). Total relief of rest pain in the iloprost group was 63% (43/68 participants) compared to 28% (18/65 participants) in the aspirin group (RR 2.28; 95% CI 1.48 to 3.52; $P = 0.0002$; [Analysis 2.3](#)).

(3) Rate of amputation: Six months after the start of treatment, three participants treated with iloprost and eight treated with aspirin required major amputation. This finding was not statistically significant (RR 0.32; 95% CI 0.09 to 1.15; $P = 0.08$; [Analysis 2.4](#)).

Secondary outcomes

Side effects: Headache, flushing, nausea, and abdominal cramps were more common in participants treated with iloprost, but according to the study authors, no participant in either group had to be withdrawn because of side-effects.

Amputation-free survival, walking distance or pain-free walking, and ankle brachial index were not assessed by [Fiessinger 1990](#).

Intravenous prostacyclin analogue versus intravenous prostaglandin analogue

Two studies assessed this comparison ([Esato 1995](#); [Ishitobi 1991](#)). [Esato 1995](#) compared intravenous prostacyclin analogue clinprost to intravenous prostaglandin analogue alprostadil. [Ishitobi 1991](#) compared intravenous prostacyclin analogue iloprost to intravenous prostaglandin analogue alprostadil.

Primary outcomes

(1) Ulcer healing: [Esato 1995](#) assessed ulcer healing at the end of treatment (4 weeks). Ulcer healing was evaluated by an assistant doctor (a medical researcher responsible for recruiting and outcome evaluation) using metric parameters (ulcer size), presence of granulation tissue and infection status. Improvement of ischaemic ulcers were seen in 70.6% (12/17 participants) in the clinprost group compared to 56.5% (13/23) in the alprostadil group (RR 1.25, 95% of CI 0.78 to 2.00; $P = 0.36$). Total recuperation was seen in 23.5% (4/17 patients) in the clinprost group compared to 8.7% (2/23 participants) in the alprostadil group (RR 2.38, 95%

of CI 0.48 to 11.7). Both outcomes were not statistically significant.

[Ishitobi 1991](#) assessed ulcer healing at the end of treatment (day 28). Ulcer healing was assessed by an assistant doctor and was evaluated with metric parameters (ulcer size), presence of granulation tissue, and presence or absence of infection. Posteriorly classified as ulcer healing improvement in Buerger's subgroups, the iloprost group reported 41% (9/22 patients) compared to 41% (11/27) in the alprostadil group (RR 1.00, 95% CI 0.51 to 1.98). This finding was not statistically significant.

Ulcer healing data (improvement of ischaemic ulcers) from [Esato 1995](#) and [Ishitobi 1991](#) were pooled and showed no significant difference between the treatment groups (RR 1.13, 95% CI 0.76 to 1.69; participants = 89; studies = 2; $P = 0.54$; $I^2 = 0\%$; [Analysis 3.1](#); [Figure 4](#); [Esato 1995](#); [Ishitobi 1991](#)).

Figure 4. Forest plot of comparison: I Prostacyclin vs Prostaglandin E1, outcome: I.1 Ulcer healing.

(2) Pain: [Esato 1995](#) assessed pain at the end of treatment (four weeks). Pain was evaluated by an assistant doctor. Fifty percent (8/16 participants) of the clinprost group were free of pain compared with 13.6% of the alprostadil group (7/22 patients; RR 1.57; 95% CI 0.72 to 3.44; $P = 0.26$); the results were not statistically significant ([Analysis 3.2](#)).

In [Ishitobi 1991](#), pain was not measured with a validated score or scale, or a quality of life questionnaire. Pain was assessed using five levels (1 = much better, 2 = better, 3 = little better, 4 = no difference, and 5 = worse). Participants in the iloprost group reported pain as: much better = 21.7% (5/23 patients), better = 34.8% (8/23 patients), little better = 17.4% (4/23 patients), no difference = 21.7% (5/23 patients), and worse = 4.4% (1/23 patients). Participants in the alprostadil group reported pain as: much better = 28.5% (8/28 patients), better = 32.1% (9/28 patients), little better = 14.3% (4/28 patients), no difference = 21.5% (6/28 patients), and worse = 3.6% (1/28 patients). [Ishitobi 1991](#) reported that this finding was not statistically significant.

(3) Rate of amputation was not appraised by [Esato 1995](#) or [Ishitobi 1991](#).

Secondary outcomes

Side effects: [Esato 1995](#) reported that five participants in the clinprost group and two participants in the alprostadil group experienced side effects. In the clinprost group, one participant developed nausea, tinnitus and vertigo in the third week of drug administration, without serious repercussion; another three participants had modification of blood tests relating to liver function, but without further repercussions after the drug administration was ceased, and one participant developed limb edema, probably not related to the clinprost use. The alprostadil group had one participant with modification in blood tests relating to liver function, but also without further repercussions after the drug administration was ceased, and another participant experienced a heat sensation in the head, only on the first day of treatment.

Side effects were not clearly specified in [Ishitobi 1991](#) for patients with Buerger's disease. Overall, 13 participants (17.3%) in the iloprost group and 11 participants (13.9%) in the alprostadil group experienced side effects, such as headache, vomiting, and flushing. Amputation-free survival, walking distance or pain-free walking, and ankle brachial index were not appraised by [Esato 1995](#) or

[Ishitobi 1991](#).

Any pharmacological agent versus placebo or any other pharmacological agent

Folic acid versus placebo

One study assessed this comparison ([Beigi 2014](#)).

Primary outcomes

(1) Pain assessment: assessed at baseline, two months and six months, using a Visual Analogue Scale (VAS), a continuous scale, ranging from 0 to 10, where a higher score means more pain. At baseline, the mean VAS score in the folic acid group was 7.07 points (SD 2.82), and the placebo group was 5.9 points (SD 2.21), resulting in MD 1.17; CI -0.66 to 3.00; $P = 0.21$; at two months, the folic acid group was 5.45 points (SD 2.75), and the placebo group was 5.75 points (SD 1.99) resulting in MD -0.30; CI -2.04 to 1.44; $P = 0.74$. After six months, the folic acid group was 3.46 points (SD 2.57), and the placebo group was 4.82 points (SD

2.46), resulting in MD -1.36; CI -3.17 to 0.45; $P = 0.14$. None of these results were statistically significant ([Analysis 4.1](#); [Analysis 4.2](#); [Analysis 4.3](#)).

(2) Rate of amputation: assessed at baseline, two months, and six months after the beginning of treatment. At baseline, five participants (5/14) in the folic acid group and four participants (4/16) in the placebo group had minor amputations; one participant (1/14) in the folic acid group and none of the participants in the placebo group had a major amputation. There was no change in the number of participants with amputations during the entire study; that is, at two months and six months, no new cases of major or minor amputation were observed. According to the study authors, the differences between the folic acid and placebo groups were not statistically significant ([Analysis 4.4](#); [Analysis 4.5](#)).

(3) Ulcer healing was not evaluated by [Beigi 2014](#).

Secondary outcomes

Side effects: none reported.

Amputation-free survival, walking distance or pain-free walking, and ankle brachial index were not assessed by [Beigi 2014](#).

ADDITIONAL SUMMARY OF FINDINGS *[Explanation]*

Intravenous prostacyclin analogue (iloprost) versus oral aspirin for treatment of Buerger’s disease						
Patient or population: patients with Buerger’s disease Settings: hospital and community Intervention: intravenous prostacyclin analogue (iloprost) Comparison: oral aspirin						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	oral aspirin	intravenous prostacyclin analogue				
Ulcer healing Follow-up: 28 days	Study population		RR 2.65 (1.15 to 6.11)	98 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	130 per 1000	346 per 1000 (150 to 797)				
Ulcer healing Follow-up: 6 months	Study population		RR 4.03 (1.24 to 13.10)	95 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	68 per 1000	275 per 1000 (86 to 893)				
Complete relief of rest pain Follow-up: 28 days	Study population		RR 2.28 (1.48 to 3.52)	133 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	277 per 1000	631 per 1000 (410 to 975)				
Rate of amputation Follow-up: 6 months	Study population		RR 0.32 (0.09 to 1.15)	95 (1 RCT)	⊕⊕⊕○ ^{1,2} moderate	
	182 per 1000	58 per 1000 (16 to 209)				

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk Ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ one single study (doubt about reproducibility of data), downgraded by one level

² conflict of interest not stated but it was not considered sufficient to downgrade the quality of evidence

Intravenous prostacyclin analogue versus intravenous prostaglandin analogue for treatment of Buerger’s disease						
Patient or population: patients with Buerger’s disease Settings: hospital and community Intervention: intravenous prostacyclin analogue (clinprost, iloprost) Comparison: intravenous prostaglandin analogue (alprostadil)						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	intra-venous prostaglandin analogue	intravenous prostacyclin analogue				
Ulcer healing Follow-up: 28 days	Study population		RR 1.13 (0.76 to 1.69)	89 (2 RCTs)	⊕○○○ ^{1,2,3,4} very low	
	480 per 1000	542 per 1000 (365 to 811)				
	Moderate					
	486 per 1000	550 per 1000 (370 to 822)				
Complete relief of pain Follow-up: 28 days	Study population		RR 1.57 (0.72 to 3.44)	38 (1 RCT)	⊕⊕○○ ^{2,3,5} low	
	318 per 1000	500 per 1000 (229 to 1000)				
Rate of amputation Follow-up: 28 days	see comment		-			Rate of amputation was not appraised by the studies in this comparison

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk Ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ adoption of 'as-treated' (per-protocol) analyses, downgraded by one level

² absence of patients' smoking history, downgraded by one level

³ conflict of interest not stated but it was not considered sufficient to downgrade the quality of evidence

⁴ small number of participants, downgraded by one level

⁵ one single study (doubt about reproducibility of data), downgraded by one level

Folic acid versus placebo for treatment of Buerger’s disease						
Patient or population: patients with Buerger’s disease Settings: community Intervention: folic acid Comparison: placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Folic acid				
Ulcer healing	see comment		-			Ulcer healing was not appraised by the study in this comparison
Pain (0 month) VAS. Scale from: 0 to 10; higher score = more pain	The mean pain in the placebo group was 5.09 points	The mean pain in the intervention group was 1.17 higher (0.66 lower to 3.00 higher)		30 (1 RCT)	⊕○○○ ¹ very low	
Pain (2 months) VAS. Scale from: 0 to 10; higher score = more pain Follow-up: 2 months	The mean pain in the placebo group was 5.75 points	The mean pain in the intervention group was 0.3 lower (2.04 lower to 1.44 higher)		30 (1 RCT)	⊕○○○ ¹ very low	
Pain (6 months) VAS. Scale from: 0 to 10; higher score = more pain Follow-up: 6 months	The mean pain in the placebo group was 4.82 points	The mean pain in the intervention group was 1.36 lower (3.17 lower to 0.45 higher)		30 (1 RCT)	⊕○○○ ¹ very low	

Change in rate of amputation (2 months) (Difference in number of amputations at start of treatment) Follow-up: 2 months	see comment	30 (1 RCT)	⊕○○○ ¹ very low	No new cases of amputations two months after start of treatment
Change in rate of amputation (6 months) (Difference in number of amputations at start of treatment) Follow-up: 6 months	see comment	30 (1 RCT)	⊕○○○ ¹ very low	No new cases of amputations six months after start of treatment
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>CI: Confidence interval; RR: Risk ratio; MD: Mean difference; VAS: Visual analogue scale</p> <p>GRADE Working Group grades of evidence High quality: Further research is very unlikely to change our confidence in the estimate of effect. Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate. Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate. Very low quality: We are very uncertain about the estimate.</p>				

¹ participants without critical ischaemia, resulting in absence of amputations and no differences in pain score in both groups, one single small study - downgraded by three levels

DISCUSSION

Summary of main results

The five included studies evaluated 602 participants and appraised the efficacy of prostacyclin or prostaglandin analogues against another drug or placebo, in patients with Buerger's disease. Folic acid was also evaluated against placebo in one recent study.

Prostacyclin analogue iloprost, intravenously administered, was effective in healing ulcers and eradicating rest pain after 28 days of treatment, when compared with aspirin. However, this evidence was restricted to a single study (133 participants), and consequently, the reproducibility of results is questionable.

Equivalent efficacy was discovered between prostacyclin (iloprost and clinprost) and prostaglandin (alprostadil) analogues as analysed by two studies, when evaluating ulcer healing and rest pain resolution after 28 days of treatment.

Oral prostacyclin analogue iloprost was not effective in healing ischaemic ulcers or eradicating rest pain when measured at the end of treatment and six months later, when compared with placebo. Evidence related to efficacy about rest pain resolution after six months of treatment with iloprost 200 mcg was inconsistent with a dose-response relationship, and the magnitude of the effect observed was low (RR 1.28; 95% CI 1.00 to 1.64); thus, the evidence is questionable.

In patients with Buerger's disease and hyperhomocysteinaemia, folic acid, when compared with placebo, did not demonstrate a protective effect against amputations, because there was an absence of amputations in both groups. Another outcome presented was pain; results found no statistically significant differences between the groups. However, the participants chosen were classed as Fontaine II; in other words, they did not have rest pain.

In most cases, treatment side effects experienced by the participants, such as headaches, flushing, or nausea, did not lead to treatment interruptions or more serious consequences.

Overall completeness and applicability of evidence

The main objective of this review was to assess the effectiveness of the pharmacological agents administered in patients with Buerger's disease at different stages of ischaemia. However, we identified only a small number of studies (five randomised controlled trials) with patients of more severe stages of the disease (ulcers, rest pain, or both). In fact, more severe cases of Buerger's disease are often treated with therapies for limb ischaemia, such as sympathectomy and pharmacological treatment, rather than limb revascularisation.

Interventions in the five studies were limited to evaluating prostaglandin and prostacyclin analogues, and folic acid, and did not evaluate other pharmacological agents such as cilostazol, clopidogrel, and pentoxifylline versus placebo or each other.

Another important issue concerns the period of treatment. This was limited to four or eight weeks in the included studies. Perhaps prolonged administration until, for example, complete healing or total relief of rest pain, could improve the success of the treatment. Evidence suggested that iloprost, given intravenously, was more effective than aspirin in healing ischaemic ulcers and eradicating rest pain. This evidence was generated with an interesting comparative i.e. aspirin; patients with severe Buerger's disease with limited treatment options are often treated as patients with atherosclerotic aetiology.

Outcomes measured in the studies were limited to assessments of ulcer healing and pain. Particularly problematic was pain evaluation. No study used a validated pain score or scale, or quality of life questionnaires, and consequently, more detailed assessments of pain was not possible. Amputation free-survival, walking distance or pain-free walking, and ankle brachial index were not assessed in any of the included studies.

Very-low quality evidence suggested that folic acid did not provide a protective effect against amputations and rest pain in patients with Buerger's disease and hyperhomocysteinaemia. Therefore, routine folic acid administration in patients with Buerger's disease and hyperhomocysteinaemia remains questionable.

Quality of the evidence

Overall, the quality of the evidence was very low to moderate, with few studies, a small number of participants and potential for biases such as randomisation and allocation concealment methods.

We summarized the quality of the evidence for the main comparisons (see also [Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#); [Summary of findings 4](#)):

(1) Oral prostacyclin analogue (iloprost) versus placebo

Classified as moderate and low quality of evidence. The evidence was obtained from a randomised controlled trial, with overall low risk of bias. However, just one study was available for this comparison and additional data about reproducibility of the outcomes were not available. In addition, evidence suggesting relief of rest pain was found with low dose of iloprost (200 mcg) but not with high dose of iloprost (400 mcg), demonstrating absence of a dose-response effect and, consequently, downgrading the quality of the evidence for relief of rest pain after six months to low.

(2) Intravenous prostacyclin analogue iloprost versus oral aspirin

Classified as moderate quality of evidence. The evidence was obtained from a randomised controlled trial, with low risk of bias, and showed a large magnitude of effect (RR > 2.0). However, just one study was available for this comparison and additional data about reproducibility of the outcomes were not available.

(3) Intravenous prostacyclin analogue versus intravenous prostaglandin analogue

Classified as low and very low quality of evidence. The evidence was obtained from two randomised controlled trials, which were double-blinded and demonstrated low heterogeneity, but the evidence was downgraded because a small number of participants was evaluated, and most importantly, the proportion of tobacco exposure between groups, and before the start of treatment was not described. In addition, 'as-treated' (per-protocol) analyses were adopted further downgrading the quality of evidence for ulcer healing.

(4) Folic acid versus placebo

Classified as very-low quality of evidence. One study with a small number of participants was available for this comparison and additional data about reproducibility of the outcomes were not available. In addition, participants without critical ischaemia were recruited into the trial, resulting in the absence of amputations and no differences in pain score in both groups.

Potential biases in the review process

The study of [Hoshino 1997](#), which assessed the efficacy of pamiogrel (antiplatelet drug) in patients with Buerger's disease, was not included in this review because a full report was not available. Thus, a potential pharmacological agent was not evaluated. Another potential bias relates to pain assessment. For the evaluation of the treatment effect on pain, we only considered absence or presence of pain in the included studies, because none of the included studies used a validated pain scale or score for appropriate assessment.

Agreements and disagreements with other studies or reviews

We did not identify any other systematic reviews about the pharmacological treatment for Buerger's disease.

AUTHORS' CONCLUSIONS

Implications for practice

There was moderate quality evidence that intravenous iloprost (prostacyclin analogue) was more effective than aspirin for eradicating rest pain and healing ischaemic ulcers in Buerger's disease, but oral iloprost was no more effective than placebo.

Low quality evidence suggested there was no difference in ulcer healing or pain relief between intravenous prostaglandin analogue alprostadil and prostacyclin (iloprost and clinprost) in severe Buerger's disease.

Very-low quality evidence suggested there was no difference in pain and rates of amputation between folic acid and placebo in patients with Buerger's disease and hyperhomocysteinaemia.

In most cases, treatment side effects such as headaches, flushing or nausea experienced by the participants were not implicated in treatment interruptions or more serious consequences.

Further high quality trials assessing the effectiveness of pharmacological agents (intravenous or oral) in patients with Buerger's disease are needed.

Implications for research

We suggest future trials investigating pharmacological treatment of Buerger's disease should incorporate the following characteristics.

- Clearly describe the methods of randomisation and concealment of allocation;
- Include participants with mild/moderate ischaemia, such as intermittent claudication;
- Evaluate outcomes, such as walking distance, pain-free walking distance and ankle brachial index;
- Compare prostacyclin versus prostaglandin analogues;
- Investigate pharmacological agents for which there currently is no RCT evidence such as pentoxifylline, cilostazol and clopidogrel;
- Assess pain utilizing validated scores/scales;
- Assess quality of life using quality of life scales;
- Evaluate time without amputation (amputation free-survival);
- Investigate folic acid reposition in patients with Buerger's disease, hyperhomocysteinaemia and critical limb ischaemia.

ACKNOWLEDGEMENTS

The review authors would like to thank Cochrane Brazil for their methodological support.

Special thanks to the Cochrane Vascular staff (Marlene Stewart, Cathryn Broderick, and Karen Welch) for all their attention and prompt support.

Thanks to Ms. Lidia Harumi Ivasa for translating the Japanese articles so competently.

REFERENCES

References to studies included in this review

Beigi 2014 *{published data only}*

Beigi AA, Hoghoughi MA, Eshaghian A, Zade AH, Masoudpour H. The role of folic acid on the hyperhomocysteinemia in the Buerger's disease (thromboangiitis obliterans). *Journal of Research in Medical Sciences* 2014;**19**:1034–7.

Esato 1995 *{published data only}*

Esato K, Yasuda K, Abe T, Hoshino S, Ishimaru S, Hori G, et al. Studies on the clinical efficacy and safety of TTC-909 in the treatment of peripheral arterial occlusive disease: a multi-center double-blind comparison with Alprostadil incorporated in lipid microspheres. *Rinsho Iyaku* 1995;**11**: 2111–2.

Fiessinger 1990 *{published data only}*

Diehm C. Multicenter-study on Iloprost in patients with thromboangiitis obliterans (Buerger's disease). *Klinische Wochenschrift* 1989;**67**:178–9.

* Fiessinger JN, Schafer M. Trial of iloprost versus aspirin treatment for critical limb ischaemia of thromboangiitis obliterans. *Lancet* 1990;**335**(8689):555–7.

Ishitobi 1991 *{published data only}*

Ishitobi K, et al. Therapeutic efficacy of iloprost, a prostacyclin analogue for chronic arterial occlusion: a double-blind comparative study with PGE1. *Rinsho to Kenkyu* 1991;**68**:1836–50.

Verstraete 1998 *{published data only}*

Verstraete M. Oral iloprost in the treatment of thromboangiitis obliterans (Buerger's disease): A double-blind, randomised, placebo-controlled trial. The European TAO Study Group. *European Journal of Vascular and Endovascular Surgery* 1998;**15**(4):300–7.

References to studies excluded from this review

Bozkurt 2006 *{published data only}*

Bozkurt AK, Koksak C, Demirbas MY, Erdogan A, Rahman A, Demirkilic U, et al. A randomized trial of intravenous iloprost (a stable prostacyclin analogue) versus lumbar sympathectomy in the management of Buerger's disease. *International Angiology* 2006;**25**:162–8.

Coscia 1972 *{published data only}*

Coscia M. Effects of nicergoline in peripheral arteriopathies of the lower limbs. Double-blind study. *Minerva Cardioangiologica* 1972;**20**(9):497–504.

He 2007 *{published data only}*

He D. The application of Simiaoyongtangjiawei in the treatment of thromboangiitis obliterans. *Chinese Journal of Information on Traditional Chinese Medicine* 2007;**14**:76.

Musial 1986 *{published data only}*

Musial J, Wilczynska M, Sladek K, Cierniewski CS, Nizankowski R, Szczeklik A. Fibrinolytic activity of prostacyclin and iloprost in patients with peripheral arterial disease. *Prostaglandins* 1986;**31**(1):61–70.

Reichert 1975 *{published data only}*

Reichert N, Shibolet S, Adar R, Gafni J. Controlled trial of propranolol in intermittent claudication. *Clinical Pharmacology and Therapeutics* 1975;**17**:612–5.

Steinorth 1967 *{published data only}*

Steinorth G. Report on the clinical testing of Lamuran (I). *Die Medizinische Welt* 1967;**35**:2025–32.

Sun 1993 *{published data only}*

Sun DX, Mu DJ, Li XY. Point injection of prostaglandin E1 on Sanyinjiao Acupuncture (SP6) for thromboangiitis obliterans pain in 24 cases. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 1993;**13**(10).

Yang 2005 *{published data only}*

Yang L, Zhou CH, Yao T. Clinical study of treating thromboangiitis obliterans with overlapping thinning chisel therapy. *Chinese Journal of Information on TCM* 2005;**12**: 73–4.

Zelikovsky 1973 *{published data only}*

Zelikovsky A, Urca I, Shaklai M, Vries A. A clinical trial with pyridinolcarbamate in the treatment of patients with obstructive diseases of the peripheral arteries. *Japanese Heart Journal* 1973;**14**:12–21.

References to studies awaiting assessment

Hoshino 1997 *{published data only}*

Hoshino S, Nakashima M, Kubo Y, Yasuda K, Abe T, Ouchi H, et al. Results of dose finding study of Pamicogrel (KBT-3022) on chronic arterial occlusive diseases. *Rinsho Iyaku* 1997;**13**:3409–36.

Additional references

Atkins 2004

Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *British Medical Journal* 2004;**328** (7454):1490–4.

Bartolo 1987

Bartolo M, Antignani PL, Todini AR, Ricci G. Buerger's disease: etiologic role of the rickettsiae?. *Journal des Maladies Vasculaires* 1987;**12**(1):82–4.

Bedenis 2014

Bedenis R, Stewart M, Cleanthis M, Robless P, Mikhailidis DP, Stansby G. Cilostazol for intermittent claudication. *Cochrane Database of Systematic Reviews* 2014, Issue 10. [DOI: 10.1002/14651858.CD003748.pub4]

Brunton 2011

Brunton LL, Chabner BA, Knollmann BC (editors). *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. 12th Edition. New York: McGraw-Hill Companies, 2011.

Buerger 1908

Buerger L. Thrombo-angiitis obliterans: a study of the vascular lesions leading to presenile spontaneous gangrene. *American Journal of Medicine* 1908;**136**:567–80.

Cachovan 1988

Cachovan M. Epidemiology and geographic distribution of the thromboangiitis obliterans [Epidemiologic und geographisches Verteilungsmuster der thromboangiitis obliterans]. *Thromboangiitis obliterans Morbus Winiwarter-Buerger*. Heidrich H. Stuttgart, 1988:31–6.

Chapman 2003

Chapman TM, Goa KL. Cilostazol: a review of its use in intermittent claudication. *American Journal of Cardiovascular Drugs* 2003;**3**(2):117–38.

Cooper 2004

Cooper LT, Tse TS, Mikhail MA, McBane RD, Stanson AW, Ballman KV. Long-term survival and amputation risk in thromboangiitis obliterans (Buerger's disease). *Journal of the American College of Cardiology* 2004;**44**(12):2410–1.

De Haro 2009

De Haro J, Florez A, Fernandez JL, Acin F. Treatment of Buerger disease (thromboangiitis obliterans) with bosentan: a case report. *BMJ Case Reports* 2009. [DOI: 10.1136/bcr.08.2008.0691]

Dindyal 2009

Dindyal S, Kyriakides C. A review of cilostazol, a phosphodiesterase inhibitor, and its role in preventing both coronary and peripheral arterial restenosis following endovascular therapy. *Recent Patents on Cardiovascular Drug Discovery* 2009;**4**(1):6–14.

Fazeli 2013

Fazeli B. Is rickettsia the key to solving the puzzle of Buerger's disease?. *Vascular* 2013;**22**(5):393–4.

FDA 1999

US Food, Drug Administration. FDA approved drug products. <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search>. DrugDetails (accessed 25 November 2015).

Fontaine 1954

Fontaine R, Kim M, Kieny R. Surgical treatment of peripheral circulation disorders [Die chirurgische Behandlung der peripheren Durchblutungsstörungen]. *Helvetica Chirurgica Acta* 1954;**21**(5/6):499–533.

Grant 1992

Grant SM, Goa KL. Iloprost. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in peripheral vascular disease, myocardial ischaemia and extracorporeal circulation procedures. *Drugs* 1992;**43**(6):889–924.

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Iwai 2005

Iwai T, Inoue Y, Umeda M, Huang Y, Kurihara N, Koike M, et al. Oral bacteria in the occluded arteries of patients with Buerger disease. *Journal of Vascular Surgery* 2005;**42**(1):107–15.

Launay 2006

Launay D, Diot E, Pasquier E, Mouthon L, Boullanger N, Fain O, et al. Bosentan for treatment of active digital ulcers in patients with systemic sclerosis. *La Presse Médicale* 2006;**35**(4 Pt 1):587–92.

Li 2008

Li X, Iwai T, Nakamura H, Inoue Y, Chen Y, Umeda M, et al. An ultrastructural study of Porphyromonas gingivalis-induced platelet aggregation. *Thrombosis Research* 2008;**122**(6):810–9.

Malecki 2009

Malecki R, Zdrojowy K, Adamiec R. Thromboangiitis obliterans in the 21st century - a new face of disease. *Atherosclerosis* 2009;**206**(2):328–34.

Matucci-Cerinic 2011

Matucci-Cerinic M, Denton CP, Furst DE, Mayes MD, Hsu VM, Carpentier P, et al. Bosentan treatment of digital ulcers related to systemic sclerosis: results from the RAPIDS-2 randomised, double-blind, placebo-controlled trial. *Annals of the Rheumatic Diseases* 2011;**70**(1):32–8.

Novo 2004

Novo S, Coppola G, Milio G. Critical limb ischemia: definition and natural history. *Current Drug Targets* 2004;**4**(3):219–25.

Olin 2000

Olin JW. Thromboangiitis obliterans (Buerger's disease). *New England Journal of Medicine* 2000;**343**(12):864–9.

Olin 2006

Olin JW, Shih A. Thromboangiitis obliterans (Buerger's disease). *Current Opinion in Rheumatology* 2006;**18**(1):18–24.

Ruffolo 2010

Ruffolo AJ, Romano M, Ciapponi A. Prostanoids for critical limb ischaemia. *Cochrane Database of Systematic Reviews* 2010, Issue 1. [DOI: 10.1002/14651858.CD006544.pub2]

Rutherford 2005

Rutherford RB. *Vascular Surgery*. 6th Edition. Vol. 1, New York: Elsevier Inc, 2005.

Safdar 2011

Safdar Z. Treatment of pulmonary arterial hypertension: the role of prostacyclin and prostaglandin analogs. *Respiratory Medicine* 2011;**105**(6):818–27.

Shionoya 1983

Shionoya S. What is Buerger's disease?. *World Journal of Surgery* 1983;**7**(4):544–51.

von Winiwarter 1879

von Winiwarter F. A peculiar form of endarteritis and endophlebitis with gangrene of the foot [Ueber eine

eigenthümliche Form von Endarteriitis und Endophlebitis mit Gangrän des Fusses]. *Archiv für Klinische Chirurgie* 1879;**23**:202–26.

Weber 1996

Weber C, Schmitt R, Birnboeck H, Hopfgartner G, van Marle SP, Peeters PAM, et al. Pharmacokinetics and pharmacodynamics of the endothelin-receptor antagonist bosentan in healthy human subjects. *Clinical Pharmacology and Therapeutics* 1996;**60**:124–37.

Weinberg 2012

Weinberg I, Jaff MR. Nonatherosclerotic arterial disorders of the lower extremities. *Circulation* 2012;**126**(2):213–22.

References to other published versions of this review

Cacione 2014

Cacione DG, Baptista-Silva JCC. Pharmacological treatment for Buerger's disease. *Cochrane Database of Systematic Reviews* 2014, Issue 3. [DOI: 10.1002/14651858.CD011033]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Beigi 2014

Methods	Study design: RCT parallel group	
Participants	Country: Iran Nº patients: 30 (14 in folic acid group and 16 in placebo group) Setting: community Mean age: 39; 42 years in folic acid group and 36 years in placebo group Gender: all male Inclusion criteria: negative history of hypertension, diabetes mellitus, ischaemic heart disease, cerebrovascular disease, collagen vascular disease or vasculitis Exclusion criteria: quote “surgical sympathectomy or any sort of vascular bypass; using aspirin, calcium channel blocker, B6 or B12; stop smoking during the study and have not compliance of being treated. Also, patients with hypercoagulative state due to inherited thrombophilia (factor V Leiden, A202120G prothrombin variant, acquired activated protein C resistance, protein C and S deficiency, antithrombin deficiency”	
Interventions	Treatment: Folic acid group: oral 5 mg folic acid tablet (by Jallinus Pharmacy, Tehran, Iran) Placebo group: oral placebo with the same colour, size, weight and box (by Amin Pharmacy, Isfahan, Iran) Duration of treatment: 1 day - a single dose of folic acid or placebo Follow-up: 6 months	
Outcomes	Primary: Major and minor amputations	
Notes	Conflict of Interest: none declared The study authors reported the following regarding a link between homocysteine levels and Buergers disease: quote “At the beginning of the study, homocysteine level was higher than normal in 19 patients (63%). There was a significant decrease in homocysteine level during 6 months in folic acid group (P < 0.001), but there was no change in the placebo group.”	
Risk of bias		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerised randomisation
Allocation concealment (selection bias)	Low risk	There was a person responsible for allocation concealment; codified using computerised randomisation, only revealed at the end of the trial

Beigi 2014 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Doctor responsible for outcome assessment assessed the patients at the start and at the end of treatment, but was not responsible for drug administration
Incomplete outcome data (attrition bias) All outcomes	Low risk	Study without losses to follow-up
Selective reporting (reporting bias)	High risk	Did not describe the development of rest pain or ischaemic ulcers after the treatment
Other bias	High risk	Participants without critical ischaemia were eligible, resulting in low chances to be amputated. Conflict of interest: none declared

Esato 1995

Methods	Study design: RCT parallel group
Participants	Country: Japan N° patients: 135, 46 with Buerger's disease (21 in clinprost group and 25 in alprostadil group) Setting: not stated Mean age: 61.5 years Gender: not described Inclusion criteria: patients with Buerger's disease and ischaemic ulcer < 1 year or rest pain Exclusion criteria: patients with previous lower limb revascularization surgery or sympathectomy; contra-indications for prostaglandin use
Interventions	Treatment: Clinprost group: Lipidious emulsion of TTC-909 (clinprost, a prostacyclin analogue): 2 ampoules (1 ampoule with TTC-909 with 1 mL plus 1 ampoule of Lipo PGE1 placebo with 2 mL) completed with saline solution, resulting in 10 mL of solution applied daily, a total dose of 2 mcg, intravenously Alprostadil group: Lipidious emulsion with PGE 1 (alprostadil), 2 ampoules (1 ampoule with Lipo PGE1 with 2 mL plus 1 ampoule of TTC-909 placebo with 1 mL) completed with saline solution, resulting in 10 mL of solution applied daily, a total dose of 10 mcg, intravenously Duration of treatment: 4 weeks No follow-up
Outcomes	Primary: improvement of rest pain and ischaemic ulcer Secondary: safety

Esato 1995 (Continued)

Notes	Conflict of interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Stated, but did not describe methods
Allocation concealment (selection bias)	Unclear risk	Information not described
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Doctor responsible for outcome assessment assessed the patients at the start and at the end of treatment, but was not responsible for drug administration
Incomplete outcome data (attrition bias) All outcomes	High risk	Adoption of “as-treated” (per protocol) analyses
Selective reporting (reporting bias)	Low risk	All outcomes reported
Other bias	High risk	Absence of patient's smoking history

Fiessinger 1990

Methods	Study design: multicentre RCT
Participants	Country: Multicentre in Europe: (9 countries, 26 centres) Setting: in hospital and community N° patients: 152 (19 did not fulfil criteria; 133: 68 in iloprost group and 65 in aspirin group) Mean age: not stated Gender: 116 M; 36 F Inclusion criteria: patients with Buerger's disease with rest pain, ischaemic ulcers or gangrene Exclusion criteria: patients with diabetes mellitus, other inflammatory vascular diseases, and amputation or lumbar sympathectomy in the preceding 3 months Gravity of illness: ulcers or gangrene - 99 patients (65%)
Interventions	Treatment: Iloprost group: Placebo tablet identical to aspirin and a 6-hour infusion of iloprost (intravenous): in first 3 days, titration to a maximum tolerated dose or 2.0 ng/kg/min. The maximum dose at day 3 was repeated on days 4 to 28 Aspirin group: 100 mg Aspirin tablet and a 6-hour daily intravenous placebo infusion Duration of treatment: 28 days

Fiessinger 1990 (Continued)

	Follow-up: 6 months	
Outcomes	Primary: total relief of rest pain, complete healing of all trophic changes Secondary: amputation	
Notes	Conflict of Interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Low risk	Randomisation centre was utilized
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcomes assessed by a physician not involved in the patient's management, or with the study
Incomplete outcome data (attrition bias) All outcomes	Low risk	Post randomisation excluded patients were described and justified. Performed an 'intention-to-treat' analyses
Selective reporting (reporting bias)	Low risk	All outcomes assessed as described in methods. Relevant outcomes described
Other bias	Low risk	Proportion of smokers to non-smokers in the study arms at the beginning and after treatment were described

Ishitobi 1991

Methods	Study design: RCT
Participants	Country: Japan Setting: hospitalised and community Nº patients: 134 (55 with Buerger's disease; 25 in iloprost group and 30 in alprostadil group) Mean age: not specified. Gender: the proportion and numbers of participants with Buerger's disease were not specified Inclusion criteria: patients diagnosed with Buerger's disease Exclusion criteria: a) patients who had previous surgical treatment for chronic arterial obstructive disease; b) patients with necrosis, where the judgement of varicose ulcer is

	unclear; c) patients with haemorrhagic events; d) patients with serious liver, kidney, or heart disease; e) hypersensitivity to medication or drugs; e) pregnancy	
Interventions	Treatment: Iloprost group: iloprost 6 mcg in a liquid injection. One dose plus 3 doses PGE1 placebo/day; Alprostadil group (PGE1): liquid injection containing 20 mcg of alprostadil. Three doses plus 1 dose iloprost placebo/day Duration of treatment: 28 days No follow-up	
Outcomes	Primary: ulcer size, ulcer improvement Secondary: improvement of rest pain	
Notes	Conflict of interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Does not describe if the outcome assessments were done by the same person responsible for recruitment
Incomplete outcome data (attrition bias) All outcomes	High risk	More drop-out patients in the iloprost group than in the alprostadil group, actual numbers and possible reasons for drop-out not provided
Selective reporting (reporting bias)	Low risk	All outcomes reported
Other bias	High risk	Does not described the proportion of smokers to non-smokers in the groups

Verstraete 1998

Methods	Study design: Multicentre parallel RCT	
Participants	Country: six countries: Bulgaria, Poland, Portugal, Greece, France, and Israel Setting: Hospitalised during the first week of treatment; after first week, participants continued the study as out-patients Nº patients: 319 (103 in placebo group, 106 in low-dose iloprost group, and 110 in high-dose iloprost group) Mean age: 40 years Gender: 292 M; 27 F Inclusion criteria: Age under 50 years, current smoker or history of smoking, angiographic criteria compatible with Buerger’s disease with typical arteriographic findings, e. g. skip lesions below the popliteal artery, corkscrew collaterals, or both (Martorell’s sign) , or direct collaterals below the knee in the absence of atherosclerotic lesions, and history of or current superficial thrombophlebitis or vasospastic symptoms Exclusion criteria : Patients with diabetes mellitus, treated or untreated hypertension (systolic BP > 160 mmHg/diastolic BP > 95 mmHg), hypercholesterolaemia (> 260 mg/dL), atrial fibrillation (or other known causes of arterial embolism), and sympathectomy within the last 3 weeks of entering the study	
Interventions	Treatment: 3 groups Iloprost group 1: Day 1: 100 mcg (1 capsule with 50 mcg, twice daily, orally). From day 2 until the end of the study: 200 mcg (2 capsules with 50 mcg, twice daily, orally); Iloprost group 2: Day 1: 200 mcg (1 capsule with 100 mcg, twice daily, orally). From day 2 until the end of the study: 400 mcg (2 capsules with 100 mcg, twice daily, orally); Placebo group: Day 1: One capsule with iloprost placebo, twice daily, orally. From day 2 until the end of the study: two capsules with iloprost placebo, twice daily, orally Duration of treatment: 8 weeks Follow-up: 6 months after treatment	
Outcomes	Primary: total healing of most important trophic lesion Secondary: total relief of rest pain without analgesics Combined endpoint: alive without major amputation, no lesion, no rest pain, no analgesics	
Notes	Conflict of Interest was not described	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double blind

Verstraete 1998 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Does not describe whether the outcome assessments were done by the same person responsible for recruitment
Incomplete outcome data (attrition bias) All outcomes	Low risk	Losses were justified. Performed 'intention-to-treat' analyses
Selective reporting (reporting bias)	Low risk	All outcomes were described
Other bias	Unclear risk	Conflict of Interest was not described

BP: blood pressure

dL: decilitre

F: female

Kg: kilogram

Lipo PGE1: lipid emulsion of prostaglandin E1

M: male

mcg: microgram

mg: milligram

min: minute

ng: nanogram

PGE1: prostaglandin analogue E1

RCT: randomised controlled trial

TTC-909: prostacyclin analogue clinprost (isocarbacyclin methylester; methyl 5-[(1S,5S,6R,7R)-7-hydroxy-6-[(E)-(S)-3-hydroxy-1-octenyl] bicyclo[3.3.0]oct-2-en-3-yl] pentanoate)

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Bozkurt 2006	Compares pharmacological treatment versus lumbar sympathectomy
Coscia 1972	Only one patient with Buerger's disease showing improvement of claudication after use of nicergoline orally. We were unable to identify the results for this single patient with Buerger's disease
He 2007	The study is about a type of acupuncture treatment of Buerger's disease
Musial 1986	This study assessed the fibrinolytic activity of prostacyclin PGI2 and iloprost during 3 five-hour infusions on three consecutive days
Reichert 1975	Non-randomised for Buerger's disease
Steinorth 1967	No patients with Buerger's disease

(Continued)

Sun 1993	Compares a drug (prostaglandin) versus acupuncture and oral use of Rotundine/L-Tetrahydropalmatine
Yang 2005	The study is about acupuncture treatment of Buerger's disease
Zelikovsky 1973	Only two patients with Buerger's disease with no differences in ischaemic ulcer and rest pain for the combined group of participants. Results were not presented by disease classification

Characteristics of studies awaiting assessment *[ordered by study ID]*

Hoshino 1997

Methods	Controlled trial (unclear if randomised)
Participants	Patients with Buerger's disease and arteriosclerosis associated with ischaemic ulcer
Interventions	Oral pamiogrel with three different doses
Outcomes	Ulcer healing
Notes	full text not available

DATA AND ANALYSES

Comparison 1. Oral prostacyclin analogue (iloprost) versus placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Ulcer healing (8 weeks) - iloprost 200 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2 Ulcer healing (8 weeks) - iloprost 400 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3 Ulcer healing (6 months) - iloprost 200 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
4 Ulcer healing (6 months) - iloprost 400 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
5 Complete relief of rest pain (8 weeks) - iloprost 200 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6 Complete relief of rest pain (8 weeks) - iloprost 400 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
7 Complete relief of rest pain (6 months) - iloprost 200 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
8 Complete relief of rest pain (6 months) - iloprost 400 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
9 Rate of amputation - iloprost 200 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
10 Rate of amputation - iloprost 400 mcg	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Comparison 2. Intravenous prostacyclin analogue (iloprost) versus oral aspirin

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Ulcer healing (4 weeks)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2 Ulcer healing (6 months)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3 Complete relief of rest pain (4 weeks)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
4 Rate of amputation	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Comparison 3. Prostacyclin versus prostaglandin E1

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Ulcer healing	2	89	Risk Ratio (M-H, Fixed, 95% CI)	1.13 [0.76, 1.69]
2 Complete relief of pain	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Comparison 4. Folic acid versus placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain (0 month)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 Pain (2 months)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3 Pain (6 months)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
4 Change in rate of amputation (2 months)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
5 Change in rate of amputation (6 months)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected

Analysis 1.1. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 1 Ulcer healing (8 weeks) - iloprost 200 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 1 Ulcer healing (8 weeks) - iloprost 200 mcg

Analysis 1.2. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 2 Ulcer healing (8 weeks) - iloprost 400 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 2 Ulcer healing (8 weeks) - iloprost 400 mcg

Analysis 1.3. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 3 Ulcer healing (6 months) - iloprost 200 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 3 Ulcer healing (6 months) - iloprost 200 mcg

Analysis 1.4. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 4 Ulcer healing (6 months) - iloprost 400 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 4 Ulcer healing (6 months) - iloprost 400 mcg

Analysis 1.5. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 5 Complete relief of rest pain (8 weeks) - iloprost 200 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 5 Complete relief of rest pain (8 weeks) - iloprost 200 mcg

Analysis 1.6. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 6 Complete relief of rest pain (8 weeks) - iloprost 400 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 6 Complete relief of rest pain (8 weeks) - iloprost 400 mcg

Analysis 1.7. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 7 Complete relief of rest pain (6 months) - iloprost 200 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 7 Complete relief of rest pain (6 months) - iloprost 200 mcg

Analysis 1.8. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 8 Complete relief of rest pain (6 months) - iloprost 400 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 8 Complete relief of rest pain (6 months) - iloprost 400 mcg

Analysis 1.9. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 9 Rate of amputation - iloprost 200 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 9 Rate of amputation - iloprost 200 mcg

Analysis 1.10. Comparison 1 Oral prostacyclin analogue (iloprost) versus placebo, Outcome 10 Rate of amputation - iloprost 400 mcg.

Review: Pharmacological treatment for Buerger's disease

Comparison: 1 Oral prostacyclin analogue (iloprost) versus placebo

Outcome: 10 Rate of amputation - iloprost 400 mcg

Analysis 2.1. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 1 Ulcer healing (4 weeks).

Review: Pharmacological treatment for Buerger's disease

Comparison: 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin

Outcome: 1 Ulcer healing (4 weeks)

Analysis 2.2. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 2 Ulcer healing (6 months).

Review: Pharmacological treatment for Buerger's disease

Comparison: 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin

Outcome: 2 Ulcer healing (6 months)

Analysis 2.3. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 3 Complete relief of rest pain (4 weeks).

Review: Pharmacological treatment for Buerger's disease

Comparison: 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin

Outcome: 3 Complete relief of rest pain (4 weeks)

Analysis 2.4. Comparison 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin, Outcome 4 Rate of amputation.

Review: Pharmacological treatment for Buerger's disease

Comparison: 2 Intravenous prostacyclin analogue (iloprost) versus oral aspirin

Outcome: 4 Rate of amputation

Analysis 3.1. Comparison 3 Prostacyclin versus prostaglandin E1, Outcome 1 Ulcer healing.

Review: Pharmacological treatment for Buerger's disease

Comparison: 3 Prostacyclin versus prostaglandin E1

Outcome: 1 Ulcer healing

Analysis 3.2. Comparison 3 Prostacyclin versus prostaglandin E1, Outcome 2 Complete relief of pain.

Review: Pharmacological treatment for Buerger's disease

Comparison: 3 Prostacyclin versus prostaglandin E1

Outcome: 2 Complete relief of pain

Analysis 4.1. Comparison 4 Folic acid versus placebo, Outcome 1 Pain (0 month).

Review: Pharmacological treatment for Buerger's disease

Comparison: 4 Folic acid versus placebo

Outcome: 1 Pain (0 month)

Analysis 4.2. Comparison 4 Folic acid versus placebo, Outcome 2 Pain (2 months).

Review: Pharmacological treatment for Buerger's disease

Comparison: 4 Folic acid versus placebo

Outcome: 2 Pain (2 months)

Analysis 4.3. Comparison 4 Folic acid versus placebo, Outcome 3 Pain (6 months).

Review: Pharmacological treatment for Buerger's disease

Comparison: 4 Folic acid versus placebo

Outcome: 3 Pain (6 months)

Analysis 4.4. Comparison 4 Folic acid versus placebo, Outcome 4 Change in rate of amputation (2 months).

Review: Pharmacological treatment for Buerger's disease

Comparison: 4 Folic acid versus placebo

Outcome: 4 Change in rate of amputation (2 months)

Analysis 4.5. Comparison 4 Folic acid versus placebo, Outcome 5 Change in rate of amputation (6 months).

Review: Pharmacological treatment for Buerger's disease

Comparison: 4 Folic acid versus placebo

Outcome: 5 Change in rate of amputation (6 months)

APPENDICES

Appendix 1. CRS search strategy

#1	Buerger:TI,AB,KY	22
#2	Buerger*:TI,AB,KY	22
#3	MESH DESCRIPTOR Thromboangiitis Obliterans	12
#4	(thromboang* near/2 oblit*):TI,AB,KY	0
#5	(thromboang* near oblit*):TI,AB,KY	30
#6	(endangitis obliterans):TI,AB,KY	0
#7	Winiwarter:TI,AB,KY	0
#8	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7	37

Appendix 2. LILACS search strategy

(MH: "Thromboangiitis Obliterans" OR "Tromboangeítis Obliterante" OR "Tromboangeíte Obliterante" OR "Doença de Buerger" OR "C14.907.137.870" OR "C14.907.940.905") AND (DB: ("IBECS" OR "LILACS")) : 38 results

Appendix 3. ISRCTN search strategy

buerger	6
buerger*	6
thromboang*	0
(thromboang* oblit*)	0
"endagitis obliterans"	0
winiwarter	0
buerger OR buerger* OR thromboang* OR (thromboang* oblit*) OR "endagitis obliterans" OR winiwarter	6

Appendix 4. Clinicaltrials.gov search strategy

buerger	8
buerger*	0
thromboang*	0
(thromboang* oblit*)	0
“endagitis obliterans”	0
winiwarter	0
buerger OR buerger* OR thromboang* OR (thromboang* oblit*) OR “endagitis obliterans” OR winiwarter	8

Appendix 5. The Australian New Zealand Clinical Trials Registry search strategy

buerger	0
buerger*	0
thromboang*	0
(thromboang* oblit*)	0
“endagitis obliterans”	0
winiwarter	0
buerger OR buerger* OR thromboang* OR (thromboang* oblit*) OR “endagitis obliterans” OR winiwarter	0

Appendix 6. The EU clinical Trials Register search strategy

buerger	2
buerger*	2
thromboang*	0

(Continued)

(thromboang* oblit*)	0
“endagitis obliterans”	0
winiwarter	1
buerger OR buerger* OR thromboang* OR (thromboang* oblit*) OR “endagitis obliterans” OR winiwarter	21

Appendix 7. The World Health Organization International Clinical Trials Registry Platform search strategy

buerger	20
buerger*	22
thromboang*	6
(thromboang* oblit*)	6
“endagitis obliterans”	0
winiwarter	0
buerger OR buerger* OR thromboang* OR (thromboang* oblit*) OR “endagitis obliterans” OR winiwarter	24

Appendix 8. OpenGrey Database search strategy

buerger's disease	3
thromboangiitis obliterans	2
von Winiwarter disease	0
buerger's disease OR thromboangiitis obliterans OR von Wini- warter disease	3

FEEDBACK

Quality of evidence, 2 February 2016

Summary

Comment: You say that: “Compared with aspirin, intravenous prostacyclin analogue iloprost improved ulcer healing (risk ratio (RR) 2.65; 95% confidence interval (CI) 1.15 to 6.11; 98 participants; one study; moderate quality evidence), and helped to eradicate rest pain after 28 days (RR 2.28; 95% CI 1.48 to 3.52; 133 participants; one study; moderate quality evidence)”.

For ulcer healing the data are from Analysis 2.1, from a single study done 25 years ago. It showed 18/52 with iloprost and 6/46 with aspirin. The definition of moderate evidence is that “further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate”. Quite right, and here I repeated the analysis with a single misclassification error simulated, so that the numbers were 17/52 vs 7/46. The relative risk then became 2.15 (0.98 to 4.72). In other words, not significant, with results from a single patient changed.

Fiessinger 1990 has no data on 4-week ulcer healing, and so far as I can see there is no explanation from whence these numbers are derived. No is any reason given as to why the denominators in the calculations are different from both the numbers randomised (68 and 65) or those followed in the longer term (51 and 44).

Some points.

1: Fiessinger 1990 gives a beautiful explanation of a double-blind double dummy method, and while it is low risk, a simple statement of double blind is insufficient given that one treatment is an infusion and the other a tablet.

2: There is no explanation of where the numbers come from.

3: And given the very considerable literature concerning the dangers of doing sums on small numbers of patients and events, surely the true answer for ulcer healing here is that we just don't know. If one patient can make a difference between significant to not significant, the surely the quality of the evidence is very low. And that is at best. Almost all statistical results here derive from small numbers of participants and events from single studies. The reality is that we can't know given the paucity of data available.

Personally, I think that the GRADE system, with three categories where one might expect results to be altered, is the real problem. Moderate quality sounds good, but the definition of moderate that “further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate” is no different from low. very low quality is the preserve of very small numbers, as here.

Reply

The Cochrane Vascular editorial base has asked the review authors to respond to the feedback.

Contributors

Feedback: Prof Andrew Moore, University of Oxford, Cochrane author and editor

I do not have any affiliation with or involvement in any organisation with a financial interest in the subject matter of my comment

Response: Marlene Stewart, Managing Editor, Cochrane Vascular

WHAT'S NEW

Last assessed as up-to-date: 13 April 2015.

Date	Event	Description
9 February 2016	Feedback has been incorporated	Feedback received. Authors invited to respond to feedback
3 February 2016	Amended	Author order amended on request of authors
3 February 2016	New citation required but conclusions have not changed	Author order amended on request of authors

CONTRIBUTIONS OF AUTHORS

DGC is the contact person with the editorial base.

DGC drafted the clinical section of the background and will respond to the clinical comments of the referees.

DGC responded to the methodology and statistics comments of the referees.

DGC and JCCBS contributed to writing the protocol.

DGC wrote the final draft of the protocol and review.

DGC, JCCBS, CRM selected studies, extracted data, performed data analysis and wrote the review.

DGC is the guarantor of the final review.

CRM: methodological supervision.

DECLARATIONS OF INTEREST

DGC: none known

CRM: none known

JCCBS: none known

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- Chief Scientist Office, Scottish Government Health Directorates, The Scottish Government, UK.
- The Cochrane Vascular editorial base is supported by the Chief Scientist Office.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

New review author (Cristiane R Macedo) joined the review team.

INDEX TERMS

Medical Subject Headings (MeSH)

Alprostadil [therapeutic use]; Amputation [statistics & numerical data]; Aspirin [therapeutic use]; Epoprostenol [analogs & derivatives; therapeutic use]; Iloprost [therapeutic use]; Prostaglandins [therapeutic use]; Randomized Controlled Trials as Topic; Thromboangiitis Obliterans [*drug therapy; surgery]; Ulcer [drug therapy]

MeSH check words

Adult; Humans; Male; Middle Aged